arXiv:1204.5287v1 [math.AC] 24 Apr 2012

TORICNESS OF BINOMIAL EDGE IDEALS

MAHDIS SAEEDI, FARHAD RAHMATI, SEYYEDE MASOOME SEYYEDI

Abstract. Let G be a finite simple graph. In this paper we will show that the binomial edge ideal of G, J_G is toric if and only if each connected component of G is complete and in this case it is the sum of toric ideal associated to bipartite complete graphs.

1. **Introduction**

Let G be a simple graph on the vertex set $[n] = \{1, \cdots, n\}$ and K a field. Herzog, Hibi and Hreinsdotir [6] and Ohtani independently in [7] introduced binomial edge ideal J_G in $R = K[x_1, \cdots, x_n, y_1, \cdots, y_n]$ attached to G and studied their algebraic properties. They proved that J_G is a radical ideal and answered the question of when J_G is a prime ideal.

The following conditions are equivalent [7]

i) G is complete around all vertices of G.

ii) Each connected component of G is a complete graph.

iii) J_G is a prime ideal.

In this paper we answer to the question of when J_G is a lattice ideal and we will show that [theorem 3.2] J_G is a lattice ideal iff $J_G = \langle I_{G_1}, \cdots, I_{G_r} \rangle$ where I_{G_i} is the toric ideal associated to a complete bipartite graph $K_{2,n}$ equivalently J_G is a prime ideal.

2. **Preliminaries**

Let G be a simple graph and $S = K[x_1, \cdots, x_n, y_1, \cdots, y_n]$. set $f_{ij} = x_i y_j - x_j y_i$. The binomial edge ideal $J_G \subset S$ of G is the ideal generated by the binomials $f_{ij} = x_i y_j - x_j y_i$ such that $\{i, j\}$ is an edge of G.

We can see easily by definitions that:

2000 *Mathematics Subject Classification.* 13F20, 13C05 , 13C14.

Key words and phrases. lattice ideal ,toric ideal,binomial edge ideal.

Proposition 2.1.

a)Suppose that G *has an isolated vertex* 1*, and* G' *is the restriction of* G *to the vertex set* $[n] \setminus \{1\}$*,*$S' = K[x_1, \cdots, x_{n-1}, y_1, \cdots, y_{n-1}]$ *and* $J_{G'}$ *is an ideal in* S'*, then* $J_G = \langle J_{G'} \rangle$*.*

b)If G_1, G_2 *are two graphs on the vertex set* $[n]$ *then* $J_{G_1} = J_{G_2}$ *if and only if* $G_1 = G_2$*.*

c)If G_1, G_2 *are two graphs on the vertex set* $[n]$ *then*

$$J_{G_1 \cup G_2} = J_{G_1} + J_{G_2}.$$

d)If G' *is complement of* G *with respect to* K_n *then*

$$J_G + J_{G'} = J_{K_n}.$$

e)Let $G_1, \cdots, G_r$ *be the connected component of* G *then* $J_G = \langle J_{G_1}, \cdots, J_{G_r} \rangle$ *.*

Let $E(G) = \{e_1, \cdots, e_q\}$ and $f_e = x_i x_j$ where $e = \{i, j\} \in E(G)$ and $F = \{f_{e_1}, \cdots, f_{e_q}\}$. Consider the graded homomorphism of K$-$algebras:

$$\varphi : B = K[t_1, \cdots, t_q] \longrightarrow K[F],$$

induced by $\varphi(t_k) = f_{e_k}$ where $K[F]$ is the sub algebra of $K[X]$ generated by $\left\{f_{e_1}, \cdots, f_{e_q}\right\}$. The kernel of φ,I_G, is the toric ideal of $K[F]$ with respect to $f_{e_1}, \cdots, f_{e_q}$, named the toric ideal associated to G.

Let M be the $n \times q$ adjacent matrix of G. Then

$$I_G = \langle \{t^{\alpha_+} - t^{\alpha_-} | \alpha \in Z^q, M\alpha = 0\} \rangle = \langle t_1 - (x_i x_j)_1, \cdots, t_q - (x_i x_j)_q \rangle \cap B$$

This ideal I_G when G is a bipartite graph can be characterized by combinatorics properties of G. see[3]

Let $W = \{v_0, v_1, \cdots, v_r = v_0\}$ be an even cycle such that $f_i = x_{i-1} x_i$. As

$$f_1 f_3 \cdots f_{r-1} = f_2 f_4 \cdots f_r$$

the binomial

$$T_W = T_1 T_3 \cdots T_{r-1} - T_2 T_4 \cdots T_r$$

is in I_G. One says that T_W is the binomial associated to W.

Let G be a bipartite graph then I_G is generated by T_W where W is an even cycle. In particular, if G is bipartite complete graph $K_{2,n}$ with $S = \{s_1, s_2\}$,$T = \{t_1, \cdots, t_n\}$ we have

$$\varphi : S = K[x_1, \cdots, x_n, y_1, \cdots, y_n] \longrightarrow K[F]$$

induced by $\varphi(x_i) = s_1t_i$, $\varphi(y_i) = s_2t_i$, $1 \leq i \leq n$, and we have $I_{K_{2,n}} = \langle x_iy_j - x_jy_i, 1 \leq i,j \leq n, i \neq j\rangle$. So for every connected simple graph G on $[n]$,$J_G \subset I_{K_{2,n}}$ and $J_{K_n} = I_{K_{2,n}}$

By a lattice we consider a finitely generated subgroup of Z^n. A partial character ρ is a homomorphism, noted by ρ too, from a sub lattice L_ρ to the multiplicative group $K^* = K - \{0\}$.

For a partial character ρ we define $I(\rho)$ the Laurent binomial ideal in $K[X^{\pm}] = K[x_1, \cdots, x_n, x_1^{-1} \cdots, x_n^{-1}]$, generated by

$$I(\rho) = \langle x^m - \rho(m) : m \in L_\rho\rangle$$

We let $m_+, m_- \in Z_+^n$ denote the positive part and negative part of a vector $m \in Z^n$. For a partial character ρ, we define the ideal

$$I_+(\rho) = \langle x^{m_+} - \rho(m)x^{m_-} : m \in L_\rho\rangle in\ R = K[x_1, \cdots, x_n]$$

Note that $I_+(\rho) = I(\rho) \cap R$ and so $x^m - \rho(m) \in L_\rho$ if and only if $x^{m_+} - \rho(m)x^{m_-} \in I_+(\rho)$.

We call an ideal I in R a lattice ideal if there exist a partial character ρ such that $I = I_+(\rho)$

Proposition 2.2. *(Eisenbud and Sturmfels [2]) If I is a binomial ideal in $R = K[x_1, \cdots, x_n]$ not containing any monomial, then there is a unique partial character ρ such that*

$$(I : \langle x_1, \cdots, x_n\rangle^\infty) = I_+(\rho)$$

Corollary 2.3. *If I is a binomial ideal in $K[X]$ not containing any monomial and ρ is a partial character then $(I : \langle x_1, \cdots, x_n\rangle^\infty) = I(\rho)$ if and only if $I = I_+(\rho)$.*

In the other words, an ideal I is a lattice ideal if and only if there exist a partial character ρ such that $(I : \langle x_1, \cdots, x_n\rangle^\infty) = I_+(\rho)$

Corollary 2.4. *The binomial edge ideal J_G is a lattice ideal if and only if $(J_G : \langle X, Y\rangle^\infty) = J_G$*

3. Main results

Theorem 3.1. *The following conditions are equivalent:*

i) G is complete around all vertices of G.

ii) The ideal J_G is a lattice ideal.

Proof. Let J_G be a lattice ideal then by corollary 2.4 $(J_G : \langle X, Y\rangle^\infty) = J_G$. Now let $k \in V(G)$ and $i, j \in N(k)$ so $\{i, k\}, \{j, k\} \in E(G)$. We have to show that $\{i, j\} \in E(G)$ and for this we see that $x_iy_j - x_jy_i \in J_G$. since $x_iy_k - x_ky_i, x_jy_k - x_ky_j \in J_G$ and $y_k(y_ix_j - y_jx_i) = y_j(x_ky_i - x_iy_k) - y_i(x_ky_j - x_jy_k)$ so $x_iy_j - x_jy_i \in (J_G : \langle X, Y\rangle^\infty)$ and $x_iy_j - x_jy_i \in J_G$ this implies that $N(K)$ is complete for each $k \in V(G)$. Conversely, in a simple connected graph G, the ideal J_G is a lattice ideal [1] and the sum of two lattice ideal is again a lattice[2] so the proof is completed by proposition 2.1. □

Theorem 3.2. *Let $G_1, \cdots, G_r$ be the connected components of G and $|V(G_i)| = n_i$ then the following conditions are equivalent:*

i) J_G is a lattice ideal.

ii) $J_G = \langle J_{G_1}, \cdots, J_{G_r}\rangle = \sum I_{K_{2,n_i}}$ and $J_{G_i} = I_{K_{2,n_i}}$

Proof. Let J_G be a lattice ideal so by proposition 3.1 G is complete around all vertices of G. In other word each connected component of G is complete. This part of proof is completed by proposition 2.1. Conversely, let $J_G = \langle J_{G_1}, \cdots, J_{G_r}\rangle$ and $J_{G_i} = I_{K_{2,n_i}}$ that means each connected component of G is complete. So G is complete around all vertices of G and by proposition 3.1, J_G is lattice. □

Corollary 3.3. *The following conditions are equivalent:*

i) G is complete around all vertices of G.

ii) All connected components of G are complete.

iii) J_G is a prime ideal.

iv) J_G is a lattice ideal.

v) J_G is a toric ideal as a sum of toric ideals associated to some bipartite complete graphs.

Acknowledgement

The authors want to thank professor Jurgen Herzog for his useful advice.

REFERENCES

[1] E, Miller and B. Sturmfels. Combinatorial commutative algebra. Springer,(2005).

[2] D. Eisenbud and B. Sturmfels, Binomial ideals, Duke Math. J. 84, (1996) 1-45.

[3] R. H. Villarreal, *Monomial Algebras*, Monographs and Textbooks in Pureand Applied Mathematics, vol. 238, Marcel Dekker Inc, New York,(2001).

[4] Jurgen Herzog, Takayuki Hibi and Freyja HreinsdOttir , Binomial Edge Ideals, Math. AC , 2009

[5] Anargyros Katsabekis, Marcel Morales, and Apostols Thoma , Binomial Generation of The Radical of a Lattice Ideal ,Math. AC, 2010

[6] J. Herzog, T. Hibi, F. Hreinsdotir, T. Kahle, J, Rauh, Binomial edge ideals and conditional independence statements, Adv. Appl. Math. 45 (2010), 317 - 333.

[7] M. Ohtani, Graphs and Ideals generated by some 2-minors, Commun Algebra 39(3) (2011), 905917.

Faculty of Mathematics and Computer Science, Amirkabir University of Technology, P. O. Box 15875-4413 , Tehran, Iran.

E-mail address: `mahdis.saeedi@aut.ac.ir`

Faculty of Mathematics and Computer Science, Amirkabir University of Technology, P. O. Box 15875-4413 , Tehran, Iran.

E-mail address: `frahmati@aut.ac.ir`

Faculty of Mathematics and Computer Science, Amirkabir University of Technology, P. O. Box 15875-4413 , Tehran, Iran.

E-mail address: `mseyyedi@aut.ac.ir`